



ASHBURN
PEDIATRIC DENTAL CENTER
(703) 726-4333

ORTHODONTIC APPLIANCE FACT SHEET

De-Impaction Appliance

The de-impaction appliance is made of wires (and sometimes acrylic) that adapt to the top of your mouth. The wires are attached to metal bands on your baby molars. A wire with a loop on it extends back from the molar band. The doctor attaches a light elastic from this loop to a metal button that is bonded to the top of the impacted permanent tooth. This will move the tooth backwards and correct the impaction. This appliance is not removable.

Things to Know

- Avoid sticky foods such as taffy, caramels, gummy candies, starburst, skittles, and gum as they might loosen the appliance. Also, avoid hard foods such as ice, nuts, and hard candy as these might break the appliance.
- Thorough, but careful, brushing around the appliance is important to keep it and your teeth clean and free from cavities. Be gentle when brushing around the back molars so that you don't loosen the elastic.
- Initially, your teeth might feel sore after placement of this appliance. Also, you might notice extra saliva in your mouth and slight speech impairment. This is expected and should go back to normal after a few days.
- Each patient is unique, and their needs are different. For some people, the use of an over-the-counter medicine similar to that taken for a headache can be used to help the soreness. If this is needed, it is usually only for a day or two.
- Speak out loud as much as possible during the first couple of days to allow your tongue to get used to this appliance.
- Certain areas of the appliance or bands to which it attaches might bother your cheeks and tongue at first. This usually goes away after a few days, but the use of wax or wet cotton balls will help to minimize the discomfort while your mouth is adjusting. Rinsing with warm salt water or Peroxyl and the use of Benzodent will help bothered areas feel better and return to normal.
- Please call us immediately if anything feels loose or broken.

Emergencies

If you have an after-hours emergency that cannot be resolved after following the instructions provided, please call Dr. Joe on his cell phone at (703) 597-8558.